



Temprid<sup>®</sup>  
SC



Temprid<sup>®</sup>  
SC

Reg. No. L9853 Act No. 36 of 1947

**REGISTRATION HOLDER:**

Bayer (Pty.) Ltd.  
Reg. No. 1968/011192/07  
P.O. Box 143, ISANDO 1600,  
SOUTH AFRICA  
TEL: (011) 921 5911

TEMPRID<sup>®</sup> is a registered trademark of  
the Bayer Group.

Batch No.: See container  
Date manufactured: See container

UN No. 3082



84932973A / 1215



Temprid<sup>®</sup>  
SC

**INSECTICIDE**

Reg. No. L9853 Act No. 36 of 1947

A suspension concentrate with contact and systemic action,  
good knockdown and long lasting residual control.

Indoor and outdoor control of American and  
German cockroaches, flies, ants, carpet  
beetles, fishmoths, litter beetles,  
mosquitoes, fleas, ticks, hide and  
skin beetles, hide and skin larve.

**ACTIVE INGREDIENTS:**  
Imidacloprid (neonicotinoid) 251 g/l  
Beta-cyfluthrin (pyrethroid) 125 g/l

TEMPRID<sup>®</sup> is a registered  
trademark of the Bayer Group.  
UN No. 3082

**200 ml** 84968943A

Ant • Bedbug • Cockroach





**WARNINGS:**

Handle with care.  
Poisonous by contact, swallowing or inhalation.  
Toxic to fish and animals. Toxic to bees.  
May cause irritation of the skin, eyes, mucous membranes and skin sensitization.  
Keep out of reach of children, uninformed persons and animals.  
Store in a cool place away from food and feedstuffs.  
Use of this material in a manner other than in accordance with directions, may injure plants or produce undesirable results.

**PRECAUTIONS:**

Do not use in the presence of persons with known sensitivity to pyrethroids or neonicotinoids.  
In the case of signs of sensitivity, obtain medical advice.  
Do not inhale spray mist.  
Wear gloves when preparing the spray mixture.  
Wash clothing if contaminated.  
Wash immediately after accidental skin contact with soap and cold water.  
Do not eat, drink or smoke whilst applying, mixing or before washing hands and face.

1.

Prevent contamination of fish ponds, food, eating utensils and drinking water.  
Clean applicator thoroughly before re-using with other material.  
Dispose of wash water where it will not contaminate edible crops or water.  
Keep animals and children away from spilt material.  
Rinse empty container three times with a volume of water equal to 10 % of that of the container. Add the rinsings to the contents of the spray tank before destroying the container.  
Destroy empty container and do not re-use for any other purpose.

**SYMPTOMS OF POISONING**

Excitement, gastrointestinal discomfort, tremor, dizziness, headache, listlessness, nausea and vomiting, epigastric pain, muscular fasciculation of limbs, unconsciousness, convulsions and coma (very high doses).

**FIRST AID TREATMENT**

Remove patient from source of poisoning to a well-ventilated area and keep him quiet and reassured. Remove contaminated clothing and rinse contaminated body area thoroughly with plenty of soap and cold water.

2.

After eye contact, flush contamination out of eyes with clean water and call a doctor. If swallowed do not induce vomiting. Wash out mouth with water and call a doctor immediately.

**NOTE TO PHYSICIAN**

Treat symptomatically. Check blood pressure and pulse rate frequently (bradycardia and hypotonia possible). Provide supportive measures for respiratory and cardiac action. Give artificial respiration if signs of paralysis appear. Additional measures involve the elimination of the substance from the body or acceleration of its excretion (gastrolavage).

**DIRECTIONS FOR USE:** USE ONLY AS DIRECTED.

**Temprid<sup>®</sup> SC** can be used in indoor and outdoor areas of commercial and industrial premises, offices, store rooms, hospitals, schools, barracks, homes, hotels, restaurants, ships, market places, warehouses, empty silos and stables.

3.

**Shake well before measuring**

| PEST   | DOSAGE  | REMARKS   |
|--|---|---|
| Cockroaches (American and German), Fishmoths | Normal conditions: 20 ml / 10 l water<br><br>High infestation: 40 ml / 10 l water | Thoroughly spray hiding places (e.g. cracks and crevices, behind stoves and sinks, in cupboards, around drains and rubbish bins etc.) and other areas where pests occur with 50 ml spray mixture/m <sup>2</sup> . If hiding places are not directly accessible spray a barrier around them. The spray deposit should not be washed off. |

4.



| PEST         | DOSAGE   | REMARKS   |
|--------------|--|---|
| Ants         | Normal conditions:<br>20 ml /<br>10 ℓ water<br><br>High infestation:<br>40 ml / 10 ℓ water | OUTDOORS: Apply as a drench into nests or spray directly onto foraging individuals. Repeat when necessary.<br><br>INDOORS: Apply as a residual spray to surfaces regularly criss-crossed by ants with 50 ml spray mixture/m <sup>2</sup> . Repeat when necessary. |
| Bedbugs      |  | Clean all bedding and clothing. Spray hiding areas (e.g. behind pictures, cracks and crevices etc.) and bed frames.   |
| Fleas, Ticks |  | Clean or vacuum infested areas and dispose off all dust collected. Spray all infested areas, especially where pets sleep, as well as under furniture with 50 ml of spray mixture/m <sup>2</sup> . Pets should be treated with a suitable flea control product.    |

5.

| PEST                            | DOSAGE   | REMARKS   |
|---------------------------------|--|---|
| Flies, Mosquitoes               | Normal conditions:<br>20 ml /<br>10 ℓ water<br><br>High infestation:<br>40 ml / 10 ℓ water | Spray 50 ml of spray mixture/m <sup>2</sup> to areas where flying insect pests rest, such as doors, ceilings, girders, the side of animal pens, around windows and surrounding vegetation.  |
| Litter Beetle (lesser mealworm) | Normal conditions:<br>20 ml/10 ℓ water<br><br>High infestation:<br>40 ml/10 ℓ water        | Place the litter in each house after the house has been cleaned and disinfected. For best results allow disinfectant to dry before placing the litter into each house. Spray the <b>Temprid SC</b> mixture to deliver 50 ml/m <sup>2</sup> over the litter area of each house. Rake all spray treatments into the litter. |

6.

| PEST   | DOSAGE  | REMARKS   |
|--|---|---|
| Hide and Skin Beetle, Hide and Skin Larve, Carpet Beetle | Normal conditions:<br>20 ml/10 ℓ water<br><br>High infestation:<br>40 ml/10 ℓ water | Apply as a coarse spray at the rate of 50 ml/m <sup>2</sup> paying particular attention to the flesh side of the hides. |

For any enquiry, contact Bayer - **0861 022 937 (08610BAYER)**. For poisoning emergencies only, Griffon Poison Information Centre may also be contacted - 082 446 8946.

© Bayer (Pty.) Ltd. All unauthorized reproduction of this label is forbidden.

[www.bayer.co.za](http://www.bayer.co.za)

7.